

GLYCOSIDASE INHIBITORS AND METHODS OF SYNTHESIZING SAME

Patent number: WO0149674
Publication date: 2001-07-12
Inventor: PINTO BRIAN M (CA); JOHNSTON BLAIR D (CA); GHAVAMI AHMAD (CA)
Applicant: UNIV FRASER SIMON (CA); PINTO BRIAN M (CA); JOHNSTON BLAIR D (CA); GHAVAMI AHMAD (CA)
Classification:
- international: C07D333/00
- european: C07D207/12; C07D333/46; C07D345/00
Application number: WO2001CA00010 20010105
Priority number(s): US20000627434 20000728; US20000174837P 20000107

Also published as:

WO0149674 (A3)
US6455573 (B1)
CA2396688 (A1)

Cited documents:

XP002168783
XP002168784
XP002168785
XP002168786
JP11029472
more >>

[Report a data error here](#)

Abstract of WO0149674

A method for synthesizing Salacinol, its stereoisomers, and non-naturally occurring selenium and nitrogen analogues thereof having formula (I). The compounds are potentially useful as glycosidase inhibitors. The synthetic schemes comprise reacting a cyclic sulfate with a 5-membered ring sugar containing a heteroatom (X). The heteroatom preferably comprises sulfur, selenium, or nitrogen. The cyclic sulfate and ring sugar reagents may be readily prepared from carbohydrate precursors, such as D-glucose, L-glucose, D-xylose and L-xylose. The target compounds are prepared by opening of the cyclic sulfates by nucleophilic attack of the heteroatoms on the 5-membered ring sugars. The resulting heterocyclic compounds have a stable, inner salt structure comprising a heteroatom cation and a sulfate anion. The synthetic schemes yield various stereoisomers of the target compounds in moderate to good yields with limited side-reactions. In an alternative embodiment of the invention, the cyclic sulfate may be similarly reacted with a 6-membered ring sugar containing a heteroatom (X) to yield a compound having formula (XII).

Data supplied from the **esp@cenet** database - Worldwide

BEST AVAILABLE COPY

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
12 July 2001 (12.07.2001)

PCT

(10) International Publication Number
WO 01/49674 A2(51) International Patent Classification⁷: C07D 333/00

2K4 (CA). GHAVAMI, Ahmad [IR/CA]; 105 - 544 Austin Avenue, Coquitlam, British Columbia V3K 3M8 (CA).

(21) International Application Number: PCT/CA01/00010

(22) International Filing Date: 5 January 2001 (05.01.2001)

(74) Agent: BAILEY, Thomas, W.; Oyen Wiggs Green & Muttala, 480 - 601 West Cordova Street, Vancouver, British Columbia V6B 1G1 (CA).

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:

60/174,837 7 January 2000 (07.01.2000) US
09/627,434 28 July 2000 (28.07.2000) US(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(63) Related by continuation (CON) or continuation-in-part (CIP) to earlier application:

US 09/627,434 (CON)
Filed on 28 July 2000 (28.07.2000)(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).(71) Applicant (*for all designated States except US*): SIMON FRASER UNIVERSITY [CA/CA]; University/Industry Liaison Office, Room 2100, Strand Hall, Burnaby, British Columbia V5A 1S6 (CA).

Published:



— Without international search report and to be republished upon receipt of that report.

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): PINTO, Brian, M. [CA/CA]; 988 Corona Crescent, Coquitlam, British Columbia V3J 6Y9 (CA). JOHNSTON, Blair, D. [CA/CA]; 3238 West 34th Avenue, Vancouver, British Columbia V6N

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: GLYCOSIDASE INHIBITORS AND METHODS OF SYNTHESIZING SAME



(57) Abstract: A method for synthesizing Salacinol, its stereoisomers, and non-naturally occurring selenium and nitrogen analogues thereof having formula (I). The compounds are potentially useful as glycosidase inhibitors. The synthetic schemes comprise reacting a cyclic sulfate with a 5-membered ring sugar containing a heteroatom (X). The heteroatom preferably comprises sulfur, selenium, or nitrogen. The cyclic sulfate and ring sugar reagents may be readily prepared from carbohydrate precursors, such as D-glucose, L-glucose, D-xylose and L-xylose. The target compounds are prepared by opening of the cyclic sulfates by nucleophilic attack of the heteroatoms on the 5-membered ring sugars. The resulting heterocyclic compounds have a stable, inner salt structure comprising a heteroatom cation and a sulfate anion. The synthetic schemes yield various stereoisomers of the target compounds in moderate to good yields with limited side-reactions. In an alternative embodiment of the invention, the cyclic sulfate may be similarly reacted with a 6-membered ring sugar containing a heteroatom (X) to yield a compound having formula (XII).

GLYCOSIDASE INHIBITORS AND METHODS OF SYNTHESIZING SAME**Technical Field**

5


This application relates to methods for synthesizing Salacinol, its stereoisomers, and analogues thereof potentially useful as glycosidase inhibitors.

10 **Background**

In treatment of non-insulin dependent diabetes (NIDD) management of blood glucose levels is critical. One strategy for treating NIDD is to delay digestion of ingested carbohydrates, thereby lowering post-prandial blood glucose concentration. This can be achieved by administering
15 drugs which inhibit the activity of enzymes, such as glucosidases, which mediate the hydrolysis of complex starches to oligosaccharides in the small intestine. For example, carbohydrate analogues, such as acarbose, reversibly inhibit the function of pancreatic α -amylase and membrane-bound
20 intestinal α -glucoside hydrolase enzymes. In patients suffering from Type II diabetes, such enzyme inhibition results in delayed glucose absorption into the blood and a smoothing or lowering of postprandial hyperglycemia, resulting in improved glycemic control.


25 Some naturally-occurring glucosidase inhibitors have been isolated from *Salacia reticulata*, a plant native to submontane forests in Sri Lanka and parts of India (known as "Kotala himbutu" in Sinhalese). *Salacia reticulata* is a woody climbing plant which has been used in the Ayurvedic system of Indian medicine in the treatment of diabetes. Traditionally,
30 Ayurvedic medicine advised that a person suffering from diabetes should drink water left overnight in a mug carved from Kotala himbutu wood. In an article published in 1997, Yoshikawa et al. reported the isolation of the compound Salacinol from a water-soluble fraction derived from the dried roots and stems of *Salacia reticulata*.¹ Yoshikawa et al. determined the structure of

Salacinol, shown below, and demonstrated its efficacy as an α -glucosidase inhibitor.



Salacinol
(C₉H₁₈O₉S₂)

Yoshikawa et al. later reported the isolation from the roots and stems of *Salacia reticulata* of Kotalanol which was also shown to be effective as an α -glucosidase inhibitor.² Like Salacinol, Kotalanol contains a thiosugar sulfonium ion and an internal sulfate providing the counterion:




Kotalanol
(C₁₂H₂₄O₁₂S₂)


Kotalanol has been found to show more potent inhibitory activity against sucrase than Salacinol and acarbose.²

The exact mechanism of action of Salacinol and other glucosidase inhibitors has not yet been elucidated. Some known glycosidase

inhibitors, such as the indolizidine alkaloids castanospermine and swainsonine, are known to carry a positive charge at physiological pH.



Castanospermine



Swainsonine

It is believed that the mechanism of action of some known inhibitors may be at least partially explained by the establishment of stabilizing electrostatic interactions between the inhibitor and the enzyme active site carboxylate residues. It is postulated that the compounds of the present invention, which comprise positively charged sulfonium, ammonium, and selenonium ions, could function in a similar manner. It is also possible that Salacinol and other compounds of the same class may act by alteration of a transport mechanism across the intestinal wall rather than by directly binding to glucosidase enzymes.

Salacinol and Kotalanol may potentially have fewer long-term side effects than other existing oral antidiabetic agents. For example, oral administration of acarbose in the treatment of Type II diabetes results in

undesirable gastrointestinal side effects in some patients, most notably increased flatulence, diarrhoea and abdominal pain. As mentioned above, Salacinol has been used as a therapy for diabetes in the Ayurvedic system of traditional medicine for many years with no notable side effects reported.

5 Further, recent animal studies have shown that the oral ingestion of an extractive from a *Salacia reticulata* trunk at a dose of 5,000 mg/kg had no serious acute toxicity or mutagenicity in rats.³

The *Salacia reticulata* plant is, however, in relatively small

10 supply and is not readily available outside of Sri Lanka and India. Accordingly, it would be desirable if Salicinol, Kotalanol and analogues thereof could be produced synthetically.

Carbohydrate processing inhibitors have also been shown to be

15 effective in the treatment of some non-diabetic disorders, such as cancer. While normal cells display characteristic oligosaccharide structures, tumor cells display very complex structures that are usually found in embryonic tissues. It is believed that these complex structures provide signal stimuli for rapid proliferation and metastasis of tumor cells. A possible strategy for

20 therapeutic use of glucosidase inhibitors is to take advantage of the differential rates of normal vs cancer cell growth to inhibit assembly of complex oligosaccharide structures. For example, the indolizidine alkaloid swainsonine, an inhibitor of Golgi α -mannosidase II, reportedly reduces tumor cell metastasis, enhances cellular immune responses, and reduces tumor cell

25 growth in mice.⁴ Swainsonine treatment has led to significant reduction of tumor mass in human patients with advanced malignancies, and is a promising drug therapy for patients suffering from breast, liver, lung and other malignancies.^{5,6}

30 The compounds of the present invention may also find application in the treatment of Alzheimer's disease due to their stable, internal salt structure. Alzheimer's is characterized by plaque formation in the brain caused by aggregation of a peptide, β -amyloid, into fibrils. This is toxic to

neuronal cells. One can inhibit this aggregation by using detergent-like molecules. It is believed that the compounds of the present invention, which are amphipathic, may demonstrate this activity.


5 The need has therefore arisen for a new class of glycosidase inhibitors which may be synthesized in high yields from readily available starting materials and which have potential use as therapeutics.

Summary of the Invention

10

In accordance with the invention, a compound selected from the group consisting of non-naturally occurring compounds represented by the general formula (I), including stereoisomers and pharmaceutically acceptable salts thereof is disclosed,

15




20

(I)


where X is selected from the group consisting of S, Se, and NH. Such compounds include stereoisomers of Salicinol. The target compounds have a
 25 stable, internal salt structure comprising heteroatom cation X and a sulfate anion; the substituents may vary without departing from the invention. Preferably, R₁, R₂, R₃, R₄ and R₅ are the same or different and are selected from the group consisting of H, OH, SH, NH₂, halogens and constituents of compounds selected from the group consisting of cyclopropanes, epoxides,
 30 aziridines and episulfides; and R₆ is selected from the group consisting of H and optionally substituted straight chain, branched, or cyclic, saturated or unsaturated hydrocarbon radicals, such as alkyl, alkenyl, alkynyl, aryl, and alkoxy substituents containing any suitable functionality.

Processes for the production of compounds of the general formula (I) are also disclosed comprising reacting a cyclic sulfate having the general formula (II) with a 5-membered ring sugar having the general formula (III)

5






10



where X is selected from the group consisting of S, Se, and NH; R^1 and R^2 are selected from the group consisting of H and a protecting group; R^3 is selected from the group consisting of H and optionally substituted straight chain, branched, or cyclic, saturated or unsaturated hydrocarbon radicals and their protected derivatives; and R^4 , R^5 and R^6 are the same or different and are selected from the group consisting of H, OH, SH, NH_2 , halogens and constituents of compounds selected from the group consisting of cyclopropanes, epoxides, aziridines and episulfides and their protected derivatives. Preferably the cyclic sulfate is a 2,4-di-O-protected-D-or L-erythritol-1,3-cyclic sulfate, such as 2,4-O-Benzylidene-D-or L-erythritol-1,3-cyclic sulfate (i.e. R^1 and R^2 comprise a benzylidene protecting group); R^3 is H or a protected polyhydroxylated alkyl chain; and R^4 , R^5 and R^6 are selected from the group consisting of OH and a protected OH group, such as $OCH_2C_6H_5$. The synthetic processes comprise the step of opening the cyclic sulfate (II) by nucleophilic attack of the heteroatom X on the sugar (III).

In an alternative embodiment of the invention, the cyclic sulfate (II) may be reacted with a 6-membered ring sugar having the general formula (XI) to yield a compound having the general formula (XII):

30




15 where X is selected from the group consisting of S, Se and NH. In this embodiment R₁, R₂, R₃, R₄, R₅ and R₆ are the same or different and are selected from the group consisting of H, OH, SH, NH₂, halogens and constituents of compounds selected from the group consisting of cyclopropanes, epoxides, aziridines and episulfides and R₇ is selected from the group consisting of H and optionally substituted straight chain, branched, or cyclic, saturated or unsaturated hydrocarbon radicals. Preferably R¹, R² and R³ are as described above in respect of compound (II) and R⁴, R⁵, R⁶ and R⁷ are selected from the group consisting of H, OH, SH, NH₂, halogens and constituents of compounds selected from the group consisting of cyclopropanes, epoxides, aziridines and episulfides and their protected derivatives.

The application also relates to pharmaceutical compositions comprising an effective amount of a compound according to formula (I) or (XII) together with a pharmaceutically acceptable carrier and to methods of treating carbohydrate metabolic disorders, such as non-insulin dependent diabetes, or different forms of cancer or Alzheimer's disease by administering to a subject in need of such treatment an effective amount of such compounds.

Detailed Description of the Preferred Embodiments

Salacinol is a naturally occurring compound which may be extracted from the roots and stems of *Salacia reticulata*, a plant native to Sri Lanka and India. This application relates to synthetic routes for preparing Salacinol (1), and its nitrogen (2) and selenium (3) analogues shown below.



This application also relates to synthetic routes for preparing the stereoisomers of compounds (1) to (3). Such analogues and stereoisomers (including stereoisomers of Salacinol) comprise a new class of compounds which are not naturally occurring and may find use as glycosidase inhibitors.


1.0 Summary of General Synthetic Scheme

20

Scheme 1(a) below, shows the general synthetic scheme developed by the inventors for arriving at the target compounds. To synthesize different stereoisomers of Salacinol and its nitrogen and selenium analogues (A) - (C), 5-membered-ring sugars are reacted with sulfate-containing compounds in accordance with the invention (in Scheme 1(a) the letters (A), (B), and (C) represent all stereoisomers of Salacinol and its nitrogen and selenium analogues (1), (2) and (3) respectively). The inventors followed a disconnection approach for determining the preferred synthetic route. A reasonable disconnection is one that gives the 5-membered-ring sugars (D) since they can be synthesized easily from readily available carbohydrate precursors. Nucleophilic substitution at C₁ of the sulfate fragment (E) can then yield the target molecules (Scheme 1(a)). A potential problem with this approach is that the leaving group (L) might act later as a


base to abstract the acidic hydrogens of the sulfonium salt and produce unwanted products. Therefore, the cyclic sulfate (F) may be used instead of (E) to obviate the problems associated with leaving group (L). Compound (G) may similarly be used as a cyclic sulfate reagent and is a protected version of

(F).



Scheme 1(a). Disconnection approach for the synthesis of (A) - (C) (R=H, $-\text{CH}_2\text{C}_6\text{H}_5$ and L= leaving group).

Scheme 1(b) below shows generally the coupling reactions for producing the target compounds (A) - (C).



Scheme 1(b). Typical coupling reaction for the synthesis of different stereoisomers (A) - (C)


Route 1 of Scheme 1(b) shows the general strategy of reacting a cyclic sulfate with a 5-membered ring sugar to produce an intermediate compound, which may include benzyl or other protecting groups. As described in further detail below, the intermediate compound is then deprotected to yield the target compounds. The inventors have determined that Route 2 of Scheme 1(b), a possible side reaction, does not occur.

2.0 Synthesis of Reagents

Cyclic sulfates and 5-membered-ring sugars were prepared in accordance with the synthetic schemes described below. As will be apparent to a person skilled in the art, other equivalent schemes for producing the reagents of the invention could be substituted.


2.1 Cyclic sulfates

Cyclic sulfates were prepared in analogous fashion to the ethylidene acetal.⁸ The cyclic sulfate (7) was synthesized in 4 steps starting from D-glucose (Scheme 2). 2,4-O-Benzylidene-D-erythrithol (5) was synthesized from D-glucose in two steps,^{9,10} and then treated with thionyl chloride to yield the cyclic sulfite (6) which was oxidized to the cyclic sulfate (7) as described by Calvo-Flores et al.⁸



Scheme 2. Synthesis of the cyclic sulfate (7).

The enantiomer (10) was also synthesized using the same route but starting from L-glucose (Scheme 3).




10

Scheme 3. Synthesis of the cyclic sulfate (10).

2.2 Synthesis of 5-Membered-ring Heterocycles


15 In order to synthesize one of the 5-membered-ring sugars (D, X=S), 1,4-anhydro-3-O-benzyl-4-thio-D-arabinitol (11), was synthesized in 9 steps starting from D-glucose (Scheme 4).¹¹ Benzylation of the compound (11), using benzyl bromide in DMF yielded 1,4-anhydro-2,3,5-tri-O-benzyl-4-thio-D-arabinitol (12) in 90% yield. Compound (11) was debenzylated to give 20 1,4-anhydro-4-thio-D-arabinitol (13) in 97% yield using a Birch reduction.



Scheme 4. Synthesis of compounds (11) - (13).

Scheme 4. Synthesis of compounds (11) - (13).


The L-isomer, 1,4-anhydro-2,3,5-tri-O-benzyl-4-thio-L-arabinitol (14) was synthesized in 5 steps starting from D-xylose (Scheme 5).¹²



5


Scheme 5. Synthesis of compounds (14)-(17)

1,4-Di-O-methanesulfonyl-2,3,5-tri-O-benzyl-D-xylitol (15) is also a key intermediate for the synthesis of the aza and seleno sugars (16) and (17). 1,4-Dideoxy-1,4-imino-L-arabinitol (16)¹³ was synthesized in 7 steps starting from D-xylose (Scheme 5). The enantiomer (19)¹³ was synthesized in an analogous way starting from L-xylose (Scheme 6). Compound (19) was also synthesized in 10 steps starting from D-xylose.¹³ 1,4-Anhydro-2,3,5-tri-O-benzyl-4-seleno-D-arabinitol (20) was synthesized in 5 steps starting from L-xylose (Scheme 6). To synthesize compound (20), Na₂Se was made in-situ by treatment of selenium metal with sodium in liquid ammonia.



Scheme 6. Synthesis of compounds (19) and (20).

Scheme 6(a) below shows a more generalized scheme for synthesizing compound (20) using other possible protecting groups ($R = \text{COR}$, $\text{CH}_2\text{C}_6\text{H}_4\text{-OMe}_p$).




Scheme 6(a). Synthesis of compounds (19) and (20). ($R = \text{COR}$, $\text{CH}_2\text{C}_6\text{H}_4\text{-OMe}_p$).

3.0 Synthesis of the Target Compounds

The target compounds (1) – (3) were prepared by opening of the cyclic sulfates by nucleophilic attack of the heteroatoms on the 5-membered rings (Scheme 1(b) above). The heteroatom gives rise to a positively charged cation and the cyclic sulfate gives rise to a negatively charged counterion. This internal salt structure may explain the stability of the target compounds toward decomposition by further nucleophilic attack.


3.1 Synthesis of Salacinol

Salacinol (1) was synthesized by nucleophilic substitution of the protected thio-arabinitol (12) with the cyclic sulfate (10) (1.2 equiv) in dry acetone containing K_2CO_3 , to give the protected intermediate compound (21) in 33% yield. Hydrogenolysis of the benzyl and benzylidene groups in AcOH:H₂O, 4:1 afforded Salacinol (1) in 67% yield (Scheme 7).




Scheme 7. Synthesis of Salacinol (1)

The same procedure was used to prepare intermediate compound (22) in 79% yield from the enantiomeric cyclic sulfate (7). Deprotection as before gave compound (23) in 59% yield (Scheme 8). Compound (23) is a diastereomer of Salacinol (1).




Scheme 8. Synthesis of compound (23)

5 Compound (24) was prepared in 40% yield from (7) and the enantiomeric thio-ether (14) (Scheme 9). Deprotection in 80% yield gave the enantiomer of Salacinol (25).



10 **Scheme 9.** Synthesis of compound (25)

To reduce the number of synthetic steps, the inventors attempted the coupling reactions with the deprotected thio-arabinotols. Thus, the partially deprotected compound (11) was reacted with the cyclic sulfate (10) in acetone, to give compound (26) in 32% yield. Deprotection yielded
15 Salacinol (1) in 36% yield (Scheme 10).




Scheme 10. Synthesis of Salacinol (1)

5 The fully-protected thio-arabinitol (13) was not soluble in acetone and the reaction in methanol produced several products.

3.2 Synthesis of Selenium Analogues

10 The seleno-analogue intermediate (27) ($R = \text{CH}_2\text{C}_6\text{H}_5$) was made starting from the seleno-arabinitol (20) ($R = \text{CH}_2\text{C}_6\text{H}_5$) and the cyclic sulfate (10) in excellent yield 86% (Scheme 11), but NMR spectroscopy showed the presence of two isomers in a ratio of 7:1 that differed in stereochemistry at the stereogenic selenium center. The isomers were separable by analytical


15 HPLC. The inventors have assigned the name "Blintol" to the new selenium analogue (3).



$R = \text{H}, \text{COR}, \text{CH}_2\text{C}_6\text{H}_5, \text{CH}_2\text{C}_6\text{H}_4\text{-OMe}_p$

Scheme 11. Synthesis of Blintol (3)

The seleno-analogue intermediate (28) ($R=CH_2C_6H_5$) was made starting from the seleno-arabinitol (20) ($R=CH_2C_6H_5$) and the cyclic sulfate (7) in excellent yield 97% (Scheme 12); a mixture of two isomers in a ratio of 3:1 that differed in stereochemistry at the stereogenic selenium center was obtained. The isomers were separable by analytical HPLC.



$R = H, COR, CH_2C_6H_5, CH_2C_6H_4-OMe_p$

Scheme 12. Synthesis of compound (29)

10


Compound (29) is a diastereomer of Blintol (3).

3.3 Synthesis of Nitrogen Analogues

15


The nitrogen analogue intermediate (30) was made by the reaction of the deprotected imino-arabinitol (19) with the cyclic sulfate (10) in a good yield 72% (Scheme 13). Compound (19) was not soluble in acetone so the reaction was performed in dry methanol. A side product (19%) which was identified to be the product of methanolysis of the cyclic sulfate was obtained. The inventors have assigned the name "Ghavamiol" to the new nitrogen analogue (2). Compound (30) was deprotected to give Ghavamiol (2) in 64% yield.

20



Scheme 13. Synthesis of Ghavamiol (2)

5 The enantiomer intermediate (31) was made by the reaction of the deprotected imino-arabinitol (16) with the cyclic sulfate (7) in a good yield 72% (Scheme 14). A side product (21%) which was identified to be the product of methanolysis of the cyclic sulfate was obtained. Compound (31) was deprotected to give compound (32) in 77% yield. Compound (32) is the
10 enantiomer of Ghavamiol (2).



Scheme 14. Synthesis of compound (32)

15


4.0 Alternative Synthetic Scheme

In an alternative embodiment of the invention, target compounds having potential application as glycosidase inhibitors may be synthesized in
20 the manner described above using 6-membered rather than 5-membered ring heterocycles as reagents. As in the embodiments described above, the cyclic

sulfate (described above) is opened in the coupling reaction due to nucleophilic attack of the heteroatoms (i.e. X=S, Se, N) on the ring sugars. As will be apparent to a person skilled in the art, the general formulas for the 6-membered sugar reagent and resulting target compound are as shown below.

5

10



The 6-membered ring target compound shares the same internal salt structure as the 5-membered ring embodiment. The substituent groups may vary as described above without departing from the invention.

15

5.0 Examples

The following examples will further illustrate the invention in greater detail although it will be appreciated that the invention is not limited to the specific examples.

20

5.1 Experimental Methods

Optical rotations were measured at 20° C. ¹H and ¹³C NMR spectra were recorded at 400.13 and 100.6 MHz for proton and carbon respectively. All assignments were confirmed with the aid of two-dimensional ¹H, ¹H (COSYDFTP) or ¹H, ¹³C (INVBTP) experiments using standard Bruker pulse programs. MALDI-TOF mass spectra were obtained for samples dispersed in a 2,5-dihydroxybenzoic acid matrix using a Perseptive Biosystems Voyager-DE instrument. Silica gel for chromatography was Merck kieselgel 60. High resolution mass spectra were LSIMS (Fab), run on a Kratos Concept H double focussing mass spectrometer at 10000 RP.

30

5.2 Preparation of Intermediates

5.2.1 Example 1 - Preparation of Cyclic Sulfate (7) (Scheme 2)

5 Step 1 – 2,4-O-Benzylidene-D-erythritol (5).

Compound (5) was prepared from 4,6-O-benzylidene-D-glucose (4) according to standard procedures.^{9,10} Compound (5) has been mentioned by MacDonald et al.,¹⁰ without characterization, which is therefore dealt with here. Mp 138-139°C ; $[\alpha]_D -44^\circ$ (c 1.0, MeOH) ; ¹H NMR (CD₃OD): δ 7.53-7.28 (5H, m, Ar), 5.53 (1H, s, H-5), 4.2. (1H, dd, $J = 10.1, 3.6$ Hz, H-4a), 3.92 (1H, dd, $J = 12.1, 1.7$ Hz, H-1a), 3.74 (1H, dd, $J = 12.1, 5.7$ Hz, H-1b), 3.67-3.55 (3H, m, H-3, H-2, H-4b); ¹³C NMR (100.6 MHz, CD₃OD): δ 139.52 (C_{ipso}), 129.77 (C_{para}), 128.99, 127.49 (4C_{ortho+meta}), 102.36 (C-5), 84.22 (C-3), 72.21 (C-4), 62.76 (C-1), 62.59 (C-2); MALDI-TOF MS: m/e 211 (M⁺ + H), 233 (M⁺ + Na). Anal. Calcd for C₁₁H₁₄O₄: C, 62.83; H, 6.72. Found: C, 62.96; H, 6.55.

Step 2 – 2,4-O-Benzylidene-D-erythritol-1,3-cyclic sulfite (6).

A solution of the diol (5) (4.5g, 21 mmol) and Et₃N (11mL, 4equiv) in dry CH₂Cl₂ (90mL) was added dropwise to a solution of SOCl₂ (2.4mL, 1.5equiv) in dry CH₂Cl₂ (60mL), with stirring in an ice-bath under an N₂ atmosphere. Stirring was continued at 0°C, until TLC (hex:EtOAc, 4:1) showed complete disappearance of the starting material. The mixture was diluted with CH₂Cl₂ (150mL) and washed with H₂O (150mL) and brine (150mL). The organic solution was dried (Na₂SO₄) and concentrated on a rotary evaporator. The product was purified by flash chromatography [hex:EtOAc, 4:1 + 0.1% Et₃N] to give a mixture of two diastereomers (4.5g, 82%). One of the isomers was selectively recrystallized from EtOAc:hex. Mp 137-139°C ; $[\alpha]_D +32^\circ$ (c 1.0, CH₂Cl₂) ; ¹H NMR (CD₂Cl₂): δ 7.48-7.36 (5H, m, Ar), 5.68 (1H, s, H-5), 5.04 (1H, ddd, $J = 10.4, 9.5, 5.0$ Hz, H-3), 4.80 (1H, dd, $J = 10.4, 10.4$ Hz, H-1a), 4.24 (1H, dd, $J = 10.5, 5.0$ Hz, H-4e), 4.18 (1H, ddd, $J = 10.4, 9.5, 4.8$ Hz, H-2), 4.06 (1H, dd, $J = 10.4, 4.8$ Hz, H-1e), 3.89 (1H, dd, $J = 10.5, 10.4$ Hz, H-4a); ¹³C NMR (100.6 MHz, CD₂Cl₂): δ 137.14 (C_{ipso}), 129.74 (C_{para}), 128.65, 126.50 (4C_{ortho+meta}), 102.72 (C-5), 73.56 (C-2), 68.16 (C-4), 63.90 (C-3),

60.18 (C-1). Anal. Calcd for $C_{11}H_{12}O_5S$: C, 51.55; H, 4.72. Found: C, 51.80; H, 4.66.

Step 3 – 2,4-O-Benzylidene-D-erythritol-1,3-cyclic sulfate (7).

5 The cyclic sulfite (6) (3.5g, 14mmol) was dissolved in a mixture of MeCN (50mL) and CCl_4 (50mL), and $NaIO_4$ (4.1g, 1.5equiv) and $RuCl_3 \cdot H_2O$ (50mg) were added followed by H_2O (50mL). The mixture was stirred vigorously at rt until TLC (hex:EtOAc,4:1) showed complete disappearance of the starting material. The mixture was diluted with Et_2O (200mL) and washed with H_2O 10 (200mL) and brine (200mL). The organic solution was dried (Na_2SO_4) and concentrated on a rotary evaporator. The product was purified by flash chromatography [hex:EtOAc, 4:1 + 0.1% Et_3N] to yield a white solid (3.5g, 95%). A portion of the product was recrystallized from EtOAc:hex. Mp 115-125°C (dec); $[\alpha]_D +4^\circ$ (c 1.0, $CHCl_3$) ; 1H NMR (CD_2Cl_2): δ 7.48-7.37 (5H, m, Ar), 5.65 (1H, s, H-5), 4.86 (1H, ddd, $J = 10.2, 9.8, 5.0$ Hz, H-3), 4.76 (1H, dd, $J = 10.7, 10.5$ Hz, H-1a), 4.65 (1H, dd, $J = 10.5, 5.0$ Hz, H-1e), 4.44 (1H, dd, $J = 10.5, 5.0$ Hz, H-4e), 4.25 (1H, ddd, $J = 10.7, 9.8, 5.0$ Hz, H-2), 3.97 (1H, dd, $J = 10.5, 10.2$ Hz, H-4a); ^{13}C NMR (100.6 MHz, CD_2Cl_2): δ 136.32 (C_{ipso}), 130.03 (C_{para}), 128.74, 126.52 ($4C_{ortho+meta}$), 102.98 (C-5), 75.74 (C-3), 73.19 15 (C-1), 71.68 (C-2), 67.64 (C-4); MALDI-TOF MS: m/e 273 ($M^+ + H$), Anal. Calcd for $C_{11}H_{12}O_6S$: C, 48.52; H, 4.45. Found: C, 48.43; H, 4.39.

5.2.2 Example 2 - Preparation of thio-arabinitol (Scheme 4)

25 **1,4-Anhydro-2,3,5-tri-O-benzyl-4-thio-D-arabinitol(12).**

A mixture of 1,4-anhydro-3-O-benzyl-4-thio-D-arabinitol (1.0g, 4.2mmol) and 60% NaH (0.85g, 5equiv) in DMF (20mL) was stirred in an ice-bath for 1h. A solution of benzyl bromide (1.9mL, 3.8equiv) in DMF (5mL) was added and the solution was stirred at rt for 3h. The mixture was added to ice-water 30 (150mL) and extracted with Et_2O (150mL). The organic solution was dried (Na_2SO_4) and concentrated. The product was purified by flash chromatography [hex:EtOAc, 4:1] to give a syrup (1.6g, 90%). $[\alpha]_D +5^\circ$ (c 1.6, $CHCl_3$) ; 1H NMR ($CDCl_3$): δ 7.38-7.23 (15H, m, Ar), 4.64-4.45 (6H, m,

CHCl₃) ; ¹H NMR (CDCl₃): δ 7.38-7.23 (15H, m, Ar), 4.64-4.45 (6H, m, CH₂Ph), 4.19 (1H, dd, *J* = 8.9, 4.6 Hz, H-2), 4.11 (1H, dd, *J* = 7.2, 3.8 Hz, H-3), 3.69 (1H, dd, *J* = 8.8, 7.6 Hz, H-5a), 3.57 (1H, ddd, *J* = 7.5, 6.4, 3.6 Hz, H-4), 3.50 (1H, dd, *J* = 8.9, 6.3 Hz, H-5b), 3.08 (1H, dd, *J* = 11.4, 5.1 Hz, H-1a),
5 2.91 (1H, dd, *J* = 11.4, 4.6 Hz, H-1b). ¹³C NMR (100.6 MHz, CDCl₃): δ 138.16, 138.06, 137.88 (3C_{ipso}), 128.40-127.59 (15C_{Ar}), 85.08 (C-3), 85.04 (C-2), 73.01 (CH₂Ph), 72.34 (C-5), 71.85, 71.50 (2CH₂Ph), 48.99 (C-4), 33.10 (C-1). Anal. Calcd for C₂₆H₂₈O₃S: C, 74.25; H, 6.72. Found: C, 74.18; H, 6.53.

10 5.2.3 Example 3 - Preparation of seleno-arabinitol (Scheme 6)

1,4-Anhydro-2,3,5-tri-O-benzyl-4-seleno-D-arabinitol (20).

Selenium metal (1.1g, 14mmol) was added to liquid NH₃ (60mL) in a -50°C bath and small pieces of Na (0.71g) were added until a blue color appeared.
15 A small portion of selenium (20mg) was added to remove the blue color. NH₃ was removed by warming on a water bath and DMF was added and removed under high vacuum to remove the rest of NH₃. A solution of the mesylated compound (18) (7.4g, 12.7mmol) in DMF (100mL) was added and the mixture was stirred under N₂ in a 70°C bath for 3 h. The mixture was cooled and the
20 solvent was removed on high vacuum. The product was partitioned between CH₂Cl₂ (150mL) and water (50mL), and the organic solution was washed with water (50mL) and brine (50mL) and dried (MgSO₄). The product was purified by flash chromatography (hex:EtOAc, 3:1) to give a yellow oil (4.74g, 80%).
[α]_D +22° (c 1.3, CHCl₃); ¹H NMR (CDCl₃): δ 7.22-7.48 (15H, m, Ar), 4.67,
25 4.61 (2H, 2d, *J* = 11.8 Hz, CH₂Ph), 4.56, 4.48 (2H, 2d, *J* = 12.1 Hz, CH₂Ph), 4.53, 4.50 (2H, 2d, CH₂Ph), 4.22 (1H, dd, *J* = 10.1, 5.1 Hz, H-2), 4.07 (1H, dd, *J* = 4.6, 4.6 Hz, H-3), 3.85 (1H, dd, *J* = 9.2, 7.6 Hz, H-5a), 3.77 (1H, ddd, *J* = 7.5, 6.9, 4.5 Hz, H-4), 3.53 (1H, dd, *J* = 9.1, 6.8 Hz, H-5b), 3.11 (1H, dd, *J* = 10.4, 5.1 Hz, H-1a), 2.96 (1H, dd, *J* = 10.4, 5.3 Hz, H-1b). ¹³C NMR (100.6
30 MHz, CDCl₃): δ 138.24, 138.21, 138.06 (3C_{ipso}), 128.40-127.60 (15C_{Ar}), 85.93 (C-2), 85.63 (C-3), 72.96 (C-5, CH₂Ph), 72.14, 71.50 (2CH₂Ph), 42.59 (C-4), 23.96 (C-1). Anal. Calcd for C₂₆H₂₈O₃Se: C, 66.65; H, 6.03. Found: C, 66.49; H, 6.05.

5.2.4 Example 4 - General procedure for the synthesis of the protected sulfonium, selenonium and ammonium sulfates (21), (22), (24), (26), (27), (28), (30), (31) (Schemes 7 – 14).

5

The thio, aza or selenosugar (3mmol) and the cyclic sulfate (1.2equiv) were dissolved in dry acetone (in the case of (21), (22), (24), (26), (27) and (28)) or dry methanol (in the case of (30) and (31)) (0.5mL) and anhydrous K₂CO₃ (7mg) was added. The mixture was stirred in a Carius tube in an oil-bath
10 (75°C) overnight. The solvent was removed under reduced pressure and the product was purified by column chromatography.

1-((1',4'-Anhydro-2',3',5'-tri-O-benzyl-4'-thio-D-arabinitol)-4'-S-yl)-2,4-O-benzylidene-1-deoxy-L-erythritol-3-sulfate (21).

15 Column chromatography [CHCl₃:MeOH, 10:1 + 0.1% Et₃N] of the crude product gave an amorphous solid (33%). $[\alpha]_D -11.9^\circ$ (c 1.7, CH₂Cl₂); ¹H NMR (CD₂Cl₂): δ 7.49-7.12 (20H, m, Ar), 5.54 (1H, s, H-5), 4.59 (1H, ddd, $J = 9.9, 5.4, 4.5$ Hz, H-3), 4.55-4.33 (8H, m, 4CH₂Ph, H-2', H-4a, H-1a, H-3'), 4.29 (1H, dt, $J = 9.5, 3.0$ Hz, H-2), 4.25 and 4.15 (2H, 2d, $J = 11.9$ Hz, CH₂Ph),
20 4.04 (1H, m, H-1'a) 4.02-3.95 (2H, m, H-4', H-1b), 3.78 (1H, dd, $J = 10.7, 10.7$ Hz, H-4b), 3.74 (1H, dd, $J = 13.6, 3.8$ Hz, H-1'b), 3.62 (1H, dd, $J = 9.9, 8.6$ Hz, H-5'a), 3.54 (1H, dd, $J = 9.9, 7.2$ Hz, H-5'b); ¹³C NMR (100.6 MHz, CD₂Cl₂): δ 137.34, 137.24, 136.56, 136.39 (4C_{ipso}), 129.73-126.62 (20C_{Ar}), 101.95 (C-5), 83.75 (C-3'), 82.82 (C-2'), 76.80 (C-2), 73.73, 72.84, 72.52 (3CH₂Ph),
25 69.54 (C-4), 67.01 (C-5'), 66.48 (C-3), 65.27 (C-4'), 49.67 (C-1), 48.28 (C-1'); MALDI-TOF MS: m/e 693 (M⁺ + H). Anal. Calcd for C₃₇H₄₀O₉S₂: C, 64.14; H, 5.82. Found: C, 63.88; H, 5.83.

1-((1',4'-Anhydro-2',3',5'-tri-O-benzyl-4'-thio-D-arabinitol)-4'-S-yl)-2,4-O-benzylidene-1-deoxy-D-erythritol-3-sulfate (22).

30 Column chromatography [CHCl₃:MeOH, 10:1 + 0.1% Et₃N] of the crude product gave an amorphous solid (79%). $[\alpha]_D -46.9^\circ$ (c 0.65, CH₂Cl₂); ¹H NMR (CD₂Cl₂): δ 7.43-7.10 (20H, m, Ar), 5.49 (1H, s, H-5), 4.62-4.34 (11H, m,

CH₂Ph, H-3, H-4a, H-2', H-1a, H-3'), 4.30-4.21 (2H, m, H-2, H-4'), 3.96 (1H, dd, $J = 9.7, 6.2$ Hz, H-5'a), 3.90 (1H, dd, $J = 13.3, 3.4$ Hz, H-1b), 3.82 (1H, dd, $J = 9.8, 9.8$ Hz, H-5'b), 3.79-3.71 (2H, m, H-1'a, H-4b), 3.51 (1H, dd, $J = 13.2, 3.9$ Hz, H-1'b); ¹³C NMR (100.6 MHz, CD₂Cl₂): δ 137.62, 137.27, 136.48, 136.29 (4C_{ipso}), 129.80-126.56 (20C_{Ar}), 102.16 (C-5), 84.25 (C-3'), 82.56 (C-2'), 77.07 (C-2), 74.02, 72.74 (3CH₂Ph), 69.75 (C-4), 67.19 (C-5'), 66.82 (C-3), 65.76 (C-4'), 50.41 (C-1), 49.60 (C-1'); MALDI-TOF MS: m/e 693 ($M^+ + H$). Anal. Calcd for C₃₇H₄₀O₉S₂: C, 64.14; H, 5.82. Found: C, 64.16; H, 5.73.

10

1-((1',4'-Anhydro-2',3',5'-tri-O-benzyl-4'-thio-L-arabinitol)-4'-S-yl)-2,4-O-benzylidene-1-deoxy-D-erythritol-3-sulfate (24).

Column chromatography [CHCl₃:MeOH, 10:1 + 0.1% Et₃N] of the crude product gave an amorphous solid (40%). $[\alpha]_D^{+14.3^\circ}$ (c 1.4, CH₂Cl₂); ¹H NMR (CD₂Cl₂): δ 7.49-7.12 (20H, m, Ar), 5.55 (1H, s, H-5), 4.60 (1H, ddd, $J = 9.8, 5.5, 4.5$ Hz, H-3), 4.55-4.44 (5H, m, 3CH₂Ph, H-2', H-4a), 4.42 (1H, dd, $J = 13.3, 2.3$ Hz, H-1a), 4.39-4.34 (2H, m, CH₂Ph, H-3'), 4.28 (1H, dt, $J = 9.8, 2.9$ Hz, H-2), 4.24 and 4.14 (2H, 2d, $J = 11.9$ Hz, CH₂Ph), 4.10 (1H, d, $J = 13.4$ Hz, H-1'a), 3.98-3.90 (2H, m, H-4', H-1b), 3.78 (1H, dd, $J = 10.5, 10.5$ Hz, H-4b), 3.67 (1H, dd, $J = 13.4, 3.8$ Hz, H-1'b), 3.62 (1H, dd, $J = 9.9, 8.7$ Hz, H-5'a), 3.53 (1H, dd, $J = 9.9, 7.2$ Hz, H-5'b); ¹³C NMR (100.6 MHz, CD₂Cl₂): δ 137.32, 137.26, 136.48, 136.25 (4C_{ipso}), 129.79-126.64 (20C_{Ar}), 102.06 (C-5), 83.96 (C-3'), 82.74 (C-2'), 76.93 (C-2), 73.81, 72.97, 72.57 (3CH₂Ph), 69.59 (C-4), 67.07 (C-5'), 66.36 (C-3), 66.31 (C-4'), 49.96 (C-1), 48.52 (C-1'). Anal. Calcd for C₃₇H₄₀O₉S₂: C, 64.14; H, 5.82. Found: C, 64.13; H, 5.74.

25

1-((1',4'-Anhydro-3'-O-benzyl-4'-thio-D-arabinitol)-4'-S-yl)-2,4-O-benzylidene-1-deoxy-L-erythritol-3-sulfate (26).

Column chromatography [CHCl₃:MeOH, 10:1 + 0.1% Et₃N] of the crude product gave an amorphous solid (32%).; ¹H NMR (CD₂Cl₂): δ 7.49-7.26 (10H, m, Ar), 6.22 (1H, d, $J = 4.4$ Hz, 2'-OH), 5.54 (1H, s, H-5), 4.96 (1H, br-s, H-2'), 4.64 (1H, d, $J = 11.6$ Hz, CH₂Ph), 4.64-4.62 (1H, m, 5'-OH), 4.56 (1H,

30

d, $J = 11.6$ Hz, CH_2Ph), 4.54–4.48 (1H, m, H-3), 4.46 (1H, dd, $J = 10.5, 5.4$ Hz, H-4a), 4.33–4.25 (3H, m, H-3', H-2, H-1'a), 4.12 (1H, dd, $J = 13.5, 2.6$ Hz, H-1a), 4.12–4.09 (1H, m, H-4'), 4.01 (1H, dd, $J = 13.5, 3.4$ Hz, H-1b), 3.92–3.82 (2H, m, H-5'a, H-5'b), 3.78 (1H, dd, $J = 10.5, 10.1$ Hz, H-4b), 3.67 (1H, dd, $J = 13.5, 3.9$ Hz, H-1'b); ^{13}C NMR (100.6 MHz, CD_2Cl_2): δ 136.92, 136.73 (2C_{ipso}), 129.97–126.61 (10C_{Ar}), 102.32 (C-5), 88.45 (C-3'), 76.61 (C-2), 76.22 (C-2'), 72.96 (CH_2Ph), 71.24 (C-4'), 69.27 (C-4), 66.96 (C-3), 60.51 (C-5'), 52.43 (C-1'), 48.30 (C-1); MALDI-TOF MS: m/e 513 ($\text{M}^+ + \text{H}$). Anal. Calcd for $\text{C}_{23}\text{H}_{28}\text{O}_9\text{S}_2$: C, 53.89; H, 5.51. Found: C, 53.64; H, 5.34.

10

1-((1',4'-Anhydro-2',3',5'-tri-O-benzyl-4'-seleno-D-arabinitol)-4'-Se-yl)-2,4-O-benzylidene-1-deoxy-L-erythritol-3-sulfate (27).

Column chromatography [CHCl_3 :MeOH, 15:1] of the crude product gave an amorphous solid (86%). NMR showed the presence of two isomers (7:1) at the stereogenic selenium center which were separated on analytical HPLC [acetonitrile/ H_2O]. Anal. Calcd for $\text{C}_{37}\text{H}_{40}\text{O}_9\text{SSe}$: C, 59.99; H, 5.45. Found: C, 59.91; H, 5.44.

15

1-((1',4'-Anhydro-2',3',5'-tri-O-benzyl-4'-seleno-D-arabinitol)-4'-Se-yl)-2,4-O-benzylidene-1-deoxy-D-erythritol-3-sulfate (28).

Column chromatography [CHCl_3 :MeOH, 15:1] of the crude product gave an amorphous solid (96%). NMR showed the presence of two isomers (3:1) at the stereogenic selenium center which were separated on analytical HPLC [acetonitrile/ H_2O]. Anal. Calcd for $\text{C}_{37}\text{H}_{40}\text{O}_9\text{SSe}$: C, 59.99; H, 5.45. Found: C, 59.91; H, 5.37.

20

1-((1',4'-Dideoxy-1',4'-imino-D-arabinitol)-4'-N-yl)-2,4-O-benzylidene-1-deoxy-L-erythritol-3-sulfate (30).

A mixture of 1,4-Dideoxy-1,4-imino-D-arabinitol (19) (100mg, 0.7mmol) and 2,4-O-benzylidene-L-erythritol-1,3-cyclic sulfate (10) (235mg, 1.2equiv) were dissolved in dry MeOH (0.5mL) and anhydrous K_2CO_3 (15mg) was added. The mixture was stirred in a Carius tube in an oil-bath (75°C) overnight. The

30

solvent was removed under reduced pressure and column chromatography [CH₂Cl₂:MeOH, 4.5:1] of the crude product gave an amorphous solid (219mg, 72%). ¹H NMR (CD₃OD): δ 7.53-7.30 (5H, m, Ar), 5.61 (1H, s, H-5), 4.53 (1H, dd, *J* = 11.1, 5.2 Hz, H-4a), 4.25 (1H, m, H-2), 4.20 (1H, ddd, *J* = 9.8, 5.2, 4.4 Hz, H-3), 4.11 (1H, br-s, H-2'), 3.99-3.84 (4H, m, H-1a, H-3', H-5'a, H-5'b), 3.82 (1H, dd, *J* = 10.7, 9.8 Hz H-4b) 3.58 (1H, m, H-1'a), 3.55-3.42 (2H, m, H-1'b, H-4'), 3.38 (1H, m, H-1b); ¹³C NMR (100.6 MHz, CD₃OD): δ 138.72 (C_{ipso}), 130.12 (C_{para}), 129.21, 127.39 (4C_{ortho+meta}), 102.33 (C-5), 78.01 (C-4', C-3', C-2), 76.31 (C-2'), 70.29 (C-4), 69.02 (C-3), 62.64 (C-1'), 60.51 (C-5'), 58.46 (C-1); MALDI-TOF MS: *m/e* 428 (M⁺ + Na), 406 (M⁺ + H); HRMS. Calcd for C₁₆H₂₃O₉SN (M + H): 406.1179. Found: 406.1192.

1-((1',4'-Dideoxy-1',4'-imino-L-arabinitol)-4'-N-yl)-2,4-O-benzylidene-1-deoxy-D-erythritol-3-sulfate (31).

A mixture of 1,4-Dideoxy-1,4-imino-L-arabinitol (**16**) (80mg, 0.6mmol) and 2,4-O-benzylidene-D-erythritol-1,3-cyclic sulfate (**7**) (190mg, 1.2equiv) were dissolved in dry MeOH (0.5mL) and anhydrous K₂CO₃ (10mg) was added. The mixture was stirred in a Carius tube in an oil-bath (75°C) overnight. The solvent was removed under reduced pressure and column chromatography [CH₂Cl₂:MeOH, 5:1] of the crude product gave an amorphous solid (175mg, 72%). ¹H NMR (CD₃OD): δ 7.52-7.31 (5H, m, Ar), 5.62 (1H, s, H-5), 4.53 (1H, dd, *J* = 10.9, 5.2 Hz, H-4a), 4.28 (1H, m, H-2), 4.20 (1H, ddd, *J* = 9.7, 5.1, 4.6 Hz, H-3), 4.14 (1H, br-s, H-2'), 4.03 (1H, m, H-1a), 3.98-3.84 (3H, m, H-3', H-5'a, H-5'b), 3.81 (1H, dd, *J* = 10.9, 10 Hz H-4b) 3.63 (1H, m, H-1'a), 3.55-3.42 (2H, m, H-1'b, H-4'), 3.38 (1H, m, H-1b); ¹³C NMR (100.6 MHz, CD₃OD): δ 138.66 (C_{ipso}), 130.15 (C_{para}), 129.23, 127.40 (4C_{ortho+meta}), 102.34 (C-5), 77.81 (C-4'), 77.52 (C-3', C-2), 76.19 (C-2'), 70.27 (C-4), 68.92 (C-3), 62.68 (C-1'), 60.41 (C-5'), 58.61 (C-1); MALDI-TOF MS: *m/e* 428 (M⁺ + Na), 406 (M⁺ + H).

30

5.2.5 Example 5 - General procedure for the deprotection of the protected sulfonium sulfates (Schemes 7 - 10) and ammonium sulfates (Schemes 13 - 14)

- 5 The protected compound was dissolved in AcOH:H₂O, 4:1 (3mL) and stirred with Pd-C (80mg) under H₂ (52 psi). After 60h the reaction mixture was filtered through a pad of Celite, which was consequently washed with MeOH. The combined filtrates were concentrated and the residue was purified by column chromatography.

10

1-((1',4'-Anhydro-4'-thio-D-arabinitol)-4'-S-yl)-1-deoxy-L-erythritol-3-sulfate (1).

- Column chromatography [CHCl₃:MeOH:H₂O, 7:3:1] of the crude product gave an amorphous solid (67%). [α]_D +2.1° (c 0.48, MeOH); ¹H NMR (pyridine-d₅):
15 δ 5.25 (1H, ddd, *J* = 7.4, 3.8, 3.6 Hz, H-3), 5.14-5.09 (2H, m, H-3', H-2'), 5.00 (1H, m, H-2), 4.78 (1H, dd, *J* = 13.0, 4.9 Hz H-1a), 4.70 (1H, m, H-4'), 4.63 (1H, dd, *J* = 13.0, 4.0 Hz H-1b), 4.61 (1H, dd, *J* = 11.8, 3.7 Hz H-4a) 4.53 (2H, m, H-5'a, H-5'b), 4.38 (1H, dd, *J* = 11.8, 3.8 Hz H-4b), 4.32 (2H, br-s, H-1'a, H-1'b); ¹³C NMR (100.6 MHz, pyridine-d₅): δ 79.14 (C-3), 79.06 (C-3'), 78.18 (C-2'), 72.30 (C-4'), 67.44 (C-2), 62.05 (C-4), 59.98 (C-5'), 52.46 (C-1), 50.35 (C-1'). HRMS. Calcd for C₉H₁₈O₉S₂ (M + H): 335.0471. Found: 335.0481.

20

1-((1',4'-Anhydro-4'-thio-D-arabinitol)-4'-S-yl)-1-deoxy-D-erythritol-3-sulfate (23).

- 25 Column chromatography [CHCl₃:MeOH:H₂O, 7:3:1] of the crude product gave an amorphous solid (59%). [α]_D -35.6° (c 0.86, MeOH); ¹H NMR (pyridine-d₅): δ 5.19 (1H, ddd, *J* = 8.0, 4.1, 3.6 Hz, H-3), 5.17-5.12 (2H, m, H-2', H-3'), 5.00 (1H, ddd, *J* = 8.0, 5.3, 4.1 Hz, H-2), 4.83 (1H, dd, *J* = 13.0, 5.1 Hz H-1a), 4.78 (1H, m, H-4'), 4.65 (1H, dd, *J* = 11.9, 3.8 Hz H-4a), 4.64-4.57 (2H, m, H-5'a, H-5'b), 4.53 (1H, dd, *J* = 13.0, 4.1 Hz H-1b), 4.40 (1H, dd, *J* = 11.9, 3.8 Hz H-4b), 4.29 (1H, dd, *J* = 12.7, 3.9 Hz H-1'a), 4.17 (1H, dd, *J* = 12.7, 2.6 Hz H-1'b); ¹³C NMR (100.6 MHz, pyridine-d₅): δ 79.46 (C-3), 79.38 (C-3'), 78.94 (C-

30

2'), 71.94 (C-4'), 67.52 (C-2), 62.02 (C-4), 60.26 (C-5'), 52.64 (C-1), 51.01 (C-1'). HRMS. Calcd for C₉H₁₈O₉S₂ (M + H): 335.0471. Found: 335.0486.

1-((1',4'-Anhydro-4'-thio-L-arabinitol)-4'-S-yl)-1-deoxy-D-erythritol-3-sulfate (25).

Column chromatography [CHCl₃:MeOH:H₂O, 7:3:1] of the crude product gave an amorphous solid (80%). [α]_D +1.1° (c 1.5, MeOH); ¹H NMR (pyridine-d₅): δ 5.23 (1H, ddd, J = 7.4, 3.8, 3.7 Hz, H-3), 5.11 (1H, m, H-3'), 5.10 (1H, m, H-2'), 4.98 (1H, m, H-2), 4.76 (1H, dd, J = 11.7, 3.7 Hz H-1a), 4.70 (1H, m, H-4'), 4.63 (1H, dd, J = 11.7, 3.8 Hz H-1b), 4.60 (1H, dd, J = 11.8, 3.7 Hz H-4a), 4.51 (2H, m, H-5'a, H-5'b), 4.35 (1H, dd, J = 11.8, 4.0 Hz H-4b), 4.31 (2H, m, H-1'a, H-1'b); ¹³C NMR (100.6 MHz, pyridine-d₅): δ 79.38 (C-3, C-2'), 78.41 (C-3'), 72.51 (C-4'), 67.63 (C-2), 62.23 (C-4), 60.21 (C-5'), 52.60 (C-1), 50.57 (C-1'). HRMS. Calcd for C₉H₁₈O₉S₂ (M + H): 335.0471. Found: 335.0466.

1-((1',4'-Dideoxy-1',4'-imino-D-arabinitol)-4'-N-yl)-1-deoxy-L-erythritol-3-sulfate (2).

Column chromatography [CHCl₃:MeOH:H₂O, 7:3:1] of the crude product gave an amorphous solid (64%). ¹H NMR (CD₃OD): δ 4.26-4.20 (2H, m H-2, H-3), 4.15 (1H, m, H-2'), 3.98 (1H, br-s, H-3'), 3.94-3.87 (3H, m, H-5'a, H-5'b', H-4a), 3.81 (1H, dd, J = 12.0, 3.5 Hz H-4b), 3.74-3.62 (2H, m, H-1a, H-1'a), 3.49-3.42 (1H, m, H-1'b), 3.40-3.35 (1H, m, H-4'), 3.15 (1H, m, H-1b); ¹³C NMR (100.6 MHz, CD₃OD): δ 81.17 (C-3), 78.27 (C-3'), 77.86 (C-4'), 76.19 (C-2'), 68.07 (C-2), 62.57 (C-1'), 61.67 (C-4), 60.72 (C-1, C-5'). HRMS. Calcd for C₉H₁₈O₉SN (M + H): 318.0859. Found: 318.0863.

1-((1',4'-Dideoxy-1',4'-imino-L-arabinitol)-4'-N-yl)-1-deoxy-D-erythritol-3-sulfate (32).

Column chromatography [CHCl₃:MeOH:H₂O, 7:3:1] of the crude product gave an amorphous solid (77%). ¹H NMR (CD₃OD): δ 4.25 (1H, m H-2), 4.23 (1H, m, H-3), 4.16 (1H, br-s, H-2'), 3.99 (1H, br-s, H-3'), 3.94-3.87 (3H, m, H-5'a, H-5'b', H-4a), 3.81 (1H, dd, J = 12.1, 3.6 Hz H-4b), 3.77-3.64 (2H, m, H-1a, H-

1'a), 3.55-3.39 (2H, m, H-1'b, H-4'), 3.22 (1H, m, H-1b); ¹³C NMR (100.6 MHz, CD₃OD): δ 81.18 (C-3), 78.23 (C-3', C-4'), 76.10 (C-2'), 68.05 (C-2), 62.66 (C-1'), 61.88 (C-4), 60.49 (C-1, C-5'). HRMS. Calcd for C₉H₁₈O₉SN (M + H): 318.0859. Found: 318.0856.

5

As will be apparent to those skilled in the art in the light of the foregoing disclosure, many alterations and modifications are possible in the practice of this invention without departing from the spirit or scope thereof. Accordingly, the scope of the invention is to be construed in accordance with
10 the substance defined by the following claims.


References:

1. Yoshikawa, M. et al. *Tetrahedron Lett.* **1997**, 38(48), 8367 - 8370.
- 5 2. Yoshikawa, M. et al. *Chem. Pharm. Bull.* **1998**, 46(8), 1339-1340.
3. Shimoda, H. et al. *Journal of the Food Hygienic Society of Japan.* **1999**, 40(3), 198-205.
4. Goss, P.E. et al. *Clinical Cancer Res.* **1997**, 3, 1077-1086.
5. Mohla, S. et al. *Anticancer Res.* **1990**, 10, 1515-1522.
- 10 6. Goss, P.E. et al. *Cancer Res.* **1994**, 54, 1450-1457.
7. Eames, J. et al. *Tetrahedron Lett.* **1998**, 39(10), 1247-1250.
8. Calvo-Flores, F.G. et al. *J. Org. Chem.* **1997**, 62, 3944-3961.
9. Foster, A.B. et al. *J. Chem. Soc.* **1961**, 5005-5011.
10. MacDonald, D.L. et al. *J. Am. Chem. Soc.* **1956**, 78, 3720-3722.
- 15 11. Yoshimura, Y. et al. *J. Org. Chem.* **1997**, 62, 3140-3152.
12. Satoh, H. et al. *Bioorg. Med. Chem. Lett.* **1998**, 8(9), 989-992.
13. Fleet, G. et al. *Tetrahedron.* **1986**, 42, 5685-5692.

20

WHAT IS CLAIMED IS:

1. A non-naturally occurring compound selected from the group consisting of compounds represented by the general formula (I) and stereoisomers and pharmaceutically acceptable salts thereof:




(I)


where X is selected from the group consisting of S, Se and NH; R₁, R₂, R₃, R₄ and R₅ are the same or different and are selected from the group consisting of H, OH, SH, NH₂, halogens and constituents of compounds selected from the group consisting of cyclopropanes, epoxides, aziridines and episulfides; and R₆ is selected from the group consisting of H and optionally substituted straight chain, branched, or cyclic, saturated or unsaturated hydrocarbon radicals.

2. The compound as defined in claim 1, wherein R₆ is selected from the group consisting of cyclopropanes, epoxides, aziridines and episulfides.
3. The compound as defined in claim 1, wherein R₁, R₂, R₃, R₄ and R₆ are OH and R₅ is H.
4. The compound as defined in claim 1, wherein R₁, R₂, R₃, R₄ and R₅ are OH and R₆ is C₃H₇O₃.

5. A process for the production of the compound of claim 1, comprising reacting a cyclic sulfate having the general formula (II) with a 5-membered ring sugar having the general formula (III)



(II)



(III)

where X is selected from the group consisting of S, Se, and NH; R^1 and R^2 are selected from the group consisting of H and a protecting group; R^3 is selected from the group consisting of H and optionally substituted straight chain, branched, or cyclic, saturated or unsaturated hydrocarbon radicals and their protected derivatives; and R^4 , R^5 and R^6 are the same or different and are selected from the group consisting of H, OH, SH, NH_2 , halogens and constituents of compounds selected from the group consisting of cyclopropanes, epoxides, aziridines and episulfides and their protected derivatives

6. The process as defined in claim 5, wherein said cyclic sulfate is a 2,4-di-O-protected-D-or L-erythritol-1,3-cyclic sulfate.
7. The process as defined in claim 6, wherein said cyclic sulfate is 2,4-O-Benzylidene-D-or L-erythritol-1,3-cyclic sulfate.
8. The process as defined in claim 5, wherein R^3 is a protected polyhydroxylated alkyl chain.
9. The process as defined in claim 5, wherein R^4 , R^5 , and R^6 are selected from the group consisting of OH and $OCH_2C_6H_5$.

10. The process as defined in claim 5, comprising the step of opening the cyclic sulfate (II) by nucleophilic attack of the heteroatom X on the sugar (III).


11. The process as defined in claim 5, wherein the coupling reaction is carried out in a solvent selected from the group consisting of acetone and methanol.

12. The process as defined in claim 11, further comprising the step of adding a base to said solvent.

13. The process as defined in claim 12, wherein said base is K_2CO_3 .


14. A process for the production of the compound of claim 1 comprising:

(a) providing a cyclic sulfate having the general formula (II)



wherein R^1 and R^2 are H or a protecting group and R^3 is selected from the group consisting of H and optionally substituted straight chain, branched, or cyclic, saturated or unsaturated hydrocarbon radicals and their protected derivatives;

(b) providing a 5-membered sugar of the general formula (III),




(III)

wherein X is selected from the group consisting of S, Se, and NH and R^4 , R^5 and R^6 are OH or a protected OH group;


- (c) reacting the cyclic sulfate with the 5-membered ring to produce an intermediate compound having an internal salt structure comprising a positively charged heteroatom X and a negatively charged sulfate counterion; and
- (d) removing any protecting groups from said intermediate compound.

15. The process as defined in claim 14, wherein the removal of the protecting groups is performed by hydrogenolysis of said intermediate compound.


16. A method of synthesis of a compound according to claim 1 wherein X is S, and said method comprises the steps set out in any one of Schemes 7, 8, 9 or 10.




Scheme 7



Scheme 8




Scheme 9



Scheme 10


17. A method of synthesis of a compound according to claim 1 wherein X is Se and said method comprises the steps set out in Scheme 11.



R = H, COR, CH₂C₆H₅, CH₂C₆H₄-OMe_p

Scheme 11.


18. A method of synthesis of a compound according to claim 1 wherein X is Se and said method comprises the steps set out in Scheme 12.



R = H, COR, CH₂C₆H₅, CH₂C₆H₄-OMe_p


Scheme 12

19. A method of synthesis of a compound according to claim 1 wherein X is NH and said method comprises the steps set out in Scheme 13.




Scheme 13

20. A method of synthesis of a compound according to claim 1 wherein X is NH and said method comprises the steps set out in Scheme 14.




Scheme 14

21. A pharmaceutical composition comprising an effective amount of a compound according to claim 1 together with a pharmaceutically acceptable carrier.
22. A method of treating a carbohydrate metabolic disorder in an affected patient comprising the step of administering to said patient a therapeutically effective amount of a compound according to claim 1.
23. The method of claim 22, wherein said carbohydrate metabolic disorder is non-insulin dependent diabetes.
24. A method of treating tumor cell proliferation and metastasis in an affected patient comprising administering to said patient a therapeutically effective amount of a compound according to claim 1.
25. A non-naturally occurring stereoisomer of Salacinol selected from the group consisting of compounds (IV) and (V)




(IV)




(V)

26. A process for the production of the stereoisomers of Salacinol comprising reacting a cyclic sulfate having the general formula (II) with a 5-membered ring sugar having the general formula (III)



(II)




(III)


where X is S; R^1 and R^2 are selected from the group consisting of H and a protecting group; R^3 is selected from the group consisting of H and optionally substituted straight chain, branched, or cyclic, saturated or unsaturated hydrocarbon radicals and their protected derivatives; and R^4 , R^5 and R^6 are the same or different and are selected from the group consisting of H, OH, SH, NH_2 , halogens and constituents of compounds selected from the group consisting of cyclopropanes, epoxides, aziridines and episulfides and their protected derivatives.

27. The process as defined in claim 26, wherein said cyclic sulfate is 2,4-O-Benzylidene-D-or L-erythritol-1,3-cyclic sulfate.

28. A process for the production of a compound having the formula (VI) comprising reacting a cyclic sulfate selected from the group consisting of compounds having the formulas (VII) and (VIII) with a sugar compound having the formula (IX) where R = H, COR, CH₂C₆H₅, CH₂C₆H₄-OMe_p.




(VI)



(VII)




(VIII)



(IX)


29. A compound selected from the group consisting of compounds represented by the general formula (XII) and stereoisomers and pharmaceutically acceptable salts thereof:




(XII)

where X is selected from the group consisting of S, Se and NH; R_1 , R_2 , R_3 , R_4 , R_5 and R_6 are the same or different and are selected from the group consisting of H, OH, SH, NH_2 , halogens and constituents of compounds selected from the group consisting of cyclopropanes, epoxides, aziridines and episulfides; and R_7 is selected from the group consisting of H and optionally substituted straight chain, branched, or cyclic, saturated or unsaturated hydrocarbon radicals.

30. A process for the production of the compound of claim 29, comprising reacting a cyclic sulfate having the general formula (II) with a 6-membered ring sugar having the general formula (XI)



(II)



(XI)

where X is selected from the group consisting of S, Se, and NH; R^1 and R^2 are selected from the group consisting of H and a protecting group; R^3 is selected from the group consisting of H and optionally substituted straight chain, branched, or cyclic, saturated or unsaturated hydrocarbon radicals and their protected derivatives; and R^4 , R^5 , R^6 and R^7 are selected from the group consisting of H, OH, SH, NH_2 , halogens and constituents of compounds selected from the group consisting of cyclopropanes, epoxides, aziridines and episulfides and their protected derivatives.

31. The process of claim 30 wherein R^3 is a protected polyhydroxylated alkyl chain or its protected derivative.

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
12 July 2001 (12.07.2001)

PCT

(10) International Publication Number
WO 01/49674 A3

(51) International Patent Classification⁷: C07D 333/46,
345/00, 207/12, 335/00, 211/04, 211/92

2K4 (CA). GHAVAMI, Ahmad [IR/CA]; 105 - 544 Austin
Avenue, Coquitlam, British Columbia V3K 3M8 (CA).

(21) International Application Number: PCT/CA01/00010

(74) Agent: BAILEY, Thomas, W.; Oyen Wiggs Green & Mu-
tala, 480 - 601 West Cordova Street, Vancouver, British Co-
lumbia V6B 1G1 (CA).

(22) International Filing Date: 5 January 2001 (05.01.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/174,837 7 January 2000 (07.01.2000) US
09/627,434 28 July 2000 (28.07.2000) US

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ,
DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR,
HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR,
LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ,
NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM,
TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(63) Related by continuation (CON) or continuation-in-part
(CIP) to earlier application:
US 09/627,434 (CON)
Filed on 28 July 2000 (28.07.2000)

(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,
CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

(71) Applicant (*for all designated States except US*): SIMON
FRASER UNIVERSITY [CA/CA]; University/Industry
Liaison Office, Room 2100, Strand Hall, Burnaby, British
Columbia V5A 1S6 (CA).

Published:
— with international search report


(72) Inventors; and

(75) Inventors/Applicants (*for US only*): PINTO, Brian, M.
[CA/CA]; 988 Corona Crescent, Coquitlam, British Co-
lumbia V3J 6Y9 (CA). JOHNSTON, Blair, D. [CA/CA];
3238 West 34th Avenue, Vancouver, British Columbia V6N


(88) Date of publication of the international search report:
29 November 2001

*For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.*

(54) Title: GLYCOSIDASE INHIBITORS AND PREPARATION THEREOF



(I)



(II)

(57) Abstract: A method for synthesizing Salacinol, its stereoisomers, and non-naturally occurring selenium and nitrogen analogues thereof having formula (I). The compounds are potentially useful as glycosidase inhibitors. The synthetic schemes comprise reacting a cyclic sulfate with a 5-membered ring sugar containing a heteroatom (X). The heteroatom preferably comprises sulfur, selenium, or nitrogen. The cyclic sulfate and ring sugar reagents may be readily prepared from carbohydrate precursors, such as D-glucose, L-glucose, D-xylose and L-xylose. The target compounds are prepared by opening of the cyclic sulfates by nucleophilic attack of the heteroatoms on the 5-membered ring sugars. The resulting heterocyclic compounds have a stable, inner salt structure comprising a heteroatom cation and a sulfate anion. The synthetic schemes yield various stereoisomers of the target compounds in moderate to good yields with limited side-reactions. In an alternative embodiment of the invention, the cyclic sulfate may be similarly reacted with a 6-membered ring sugar containing a heteroatom (X) to yield a compound having formula (XII).



WO 01/49674 A3

INTERNATIONAL SEARCH REPORT

International Application No

PCT/CA 01/00010

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D333/46 C07D345/00 C07D207/12 C07D335/00 C07D211/04
C07D211/92

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

WPI Data, EPO-Internal, BIOSIS, PAJ, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P, X	YUASA, H. ET AL: "Synthesis of salacinol" TETRAHEDRON LETT. (2000), 41(34), 6615-6618 , XP002168783 page 6617 Scheme 2 ---	1-31
X	YOSHIKAWA, MASAYUKI ET AL: "Salacinol, potent antidiabetic principle with unique thiosugar sulfonium sulfate structure from the ayurvedic traditional medicine Salacia reticulata in Sri Lanka and India" TETRAHEDRON LETT. (1997), 38(48), 8367-8370 , XP002168784 cited in the application page 8368; figure 1 --- -/--	1,3,4, 21-25



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

& document member of the same patent family

Date of the actual completion of the international search

5 June 2001

Date of mailing of the international search report

13/07/2001

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl.
Fax: (+31-70) 340-3016

Authorized officer

Samsam Bakhtiary, M

INTERNATIONAL SEARCH REPORT

International Application No

PCT/CA 01/00010

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>DATABASE CHEMABS 'Online! CHEMICAL ABSTRACTS SERVICE, COLUMBUS, OHIO, US; MATSUDA, HISASHI ET AL: "Antidiabetic principles of natural medicines. IV. Aldose reductase and.alpha.-glucosidase inhibitors from the roots of Salacia oblonga WALL. (Celastraceae): structure of a new friedelane-type triterpene, kotalageni 16-acetate" retrieved from STN Database accession no. 132:163473 XP002168785 cited in the application abstract & CHEM. PHARM. BULL. (1999), 47(12), 1725-1729 ,</p>	<p>1,3,4, 21-25</p>
X	<p>----- DATABASE CHEMABS 'Online! CHEMICAL ABSTRACTS SERVICE, COLUMBUS, OHIO, US; YOSHIKAWA, MASAYUKI ET AL: "Antidiabetic constituents of Sri Lankan natural medicine Kotala himbutu (Salacia reticulata): absolute stereostructures of.alpha.-glucosidase inhibitors, salacinol and kotalanol, with unique thiosugar sulfonium sulfate inner salt structure" retrieved from STN Database accession no. 131:106694 XP002168786 abstract & TENNEN YUKI KAGOBUTSU TORONKAI KOEN YOSHISHU (1998), 40TH, 67-72 ,</p>	<p>1,3,4, 21-25</p>
X	<p>----- PATENT ABSTRACTS OF JAPAN vol. 1999, no. 05, 31 May 1999 (1999-05-31) & JP 11 029472 A (RES INST FOR PROD DEV), 2 February 1999 (1999-02-02) abstract ----- -/--</p>	<p>1,3,4, 21-25</p>

INTERNATIONAL SEARCH REPORT

International Application No

PCT/CA 01/00010

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>DATABASE CHEMABS 'Online! CHEMICAL ABSTRACTS SERVICE, COLUMBUS, OHIO, US; YOSHIKAWA, MASAYUKI ET AL: "Kotalanol, a potent.alpha.-glucosidase inhibitor with thiosugar sulfoniu sulfate structure, from antidiabetic Ayurvedic medicine salacia reticulat" retrieved from STN Database accession no. 129:300080 XP002168787 cited in the application abstract & CHEM. PHARM. BULL. (1998), 46(8), 1339-1340 ,</p> <p>-----</p>	<p>1,3,4, 21-25</p>

FURTHER INFORMATION CONTINUED FROM PCT/SA/ 210

Continuation of Box I.1

Although claims 22,23,24 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.

Continuation of Box I.1

Claims Nos.: 22,23,24

Rule 39.1(iv) PCT - Method for treatment of the human or animal body by therapy

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/CA 01/00010

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
JP 11029472 A	02-02-1999	JP 3030008 B	10-04-2000